

RECOFAST ENTERICUM

Oral powder

CUT YOUR LOSSES

Dietetic complementary feed

Composition

Rice flour, 10%; Algae extract, 7.5%; Milk serum powder, 5%; Maize starch, 5%; Dried beer yeast, 4%; Ground carob, 3%; Carrot flour, 2.5%; Sodium bicarbonate, 2.5%; Sodium chloride, 2.5%; Magnesium chloride hexahydrate, 1%; Pectin, 1%.

Analytical constituents: Crude fibre, 20-25%; Crude protein, 6-11%; Crude ash, 7-10%; Crude oils and fats, < 5%; Chlorides, 2-3%; Calcium, 1-2%; Sodium, 1-2%; Potassium, 0%; Lysine, 0%; Methionine, 0%; Water content, < 10%.

Carbohydrate sources: Rice flour, algae extract, maize starch, dried beer yeast, ground carob, and carrot flour.

Bicarbonates: Sodium bicarbonate.

Additives: Vitamin A, 135,000 IU/kg; Vitamin D₃, 20,000 IU/kg; Nicotinamide, 2.5 g/kg; Sodium ascorbate, 2.5 g/kg; Vitamin B₁ HCl, 1.25 g/kg; Vitamin E (α-tocopherol), 0.75 g/kg; Vitamin B₂ sodium phosphate, 0.5 g/kg; Calcium pantothenate, 0.5 g/kg; Vitamin B₁₂, 2.5 mg/kg; Lecithins, 50 g/kg.

Properties

Stabilization of water and electrolyte balance to support the physiological digestion. In case of risk of, during periods of, or recovery from digestive disturbance (diarrhoea).

Indications and target species

Calves, pigs, lambs, kids and foals: In case of risk of, during periods of, or recovery from digestive disturbance (diarrhoea).

Contraindications, adverse reactions and interactions

Not described.

Dosage for each species, route and method of administration

Oral use.

Calves and foals: 100 g per day, administered once daily or divided into two administrations. Administer from 1 to 7 days (1 to 3 days if fed exclusively). Dilute sachet in 100ml-1L of warm water or milk.

Lambs, kids and pigs: 20 g per day, administered once daily or divided into two administrations. Administer from 1 to 7 days (1 to 3 days if fed exclusively). Dilute 10 g in 20 ml of warm water or milk. It is recommended that advice from a veterinarian be sought before use and before extending the period of use.

Withdrawal period

Not necessary.

Storage precautions

Keep out of the sight and reach of children. Protect from moisture, light and heat.

Other information

100g sachets.

For animal use.



RECOFAST ENTERICUM

CUT YOUR LOSSES

Oral powder

Along with you

RECOFAST ENTERICUM


Oral powder

DIARRHOEA - A CONSTANT CHALLENGE TO CALF HEALTH

Diarrhoea is one of the most common causes of death in neonatal animals and is responsible for significant economic losses on farms every year.

Most cases of serious scour are caused by infectious agents such as Rotavirus/Coronavirus, Cryptosporidium and less frequently, *E.coli* and Salmonella.

Adverse management factors strongly predispose to the development of scour and worsen the outcomes from the condition.



Economic losses from scour arise from:

- Loss of the animal/carcass disposal costs
- Veterinary treatment costs
- Labour costs – animals require intensive support

ACTION PLAN FOR CALF SCOUR

- 1 Take samples from acute cases for analysis to aid future prevention strategy.
- 2 Isolate sick calves from healthy pen-mates to prevent spread.
- 3 Ensure calves are well bedded and warm.
- 4 Continue to feed milk or allow to suckle, to maintain strength and aid recovery.
- 5 If calf is dehydrated give supplemental electrolyte solution.
- 6 Administer RECOFAST to help the calf form a more normal stool, slowing water loss.



CUT YOUR LOSSES



RECOFAST ENTERICUM is a **supportive product** for animals recovering from **diarrhoea (scour)**.

FEATURES OF RECOFAST



Astringent

RECOFAST's main action is to absorb and hold water in the bowel, forming a more normal stool and slowing water loss.



Comfort

RECOFAST helps reduce the frequency of purging episodes, unproductive straining and scalding of hindquarters, thus helping relieve the discomfort due to scour.




Nutritional supplement


RECOFAST contains electrolytes, vitamins, proteins and carbohydrates.

DOSE

It can be administered once daily or divided in two administrations.



Calves and foals
100 g/day, 1-3 days.



Lambs, goat kids and piglets
20 g/day, 1-3 days.

Rapid results
No withdrawal time
Easy to use

PRACTICAL RECOMMENDATIONS

RECOFAST ENTERICUM should be administered mixed with liquid.



How to dissolve the product

RECOFAST ENTERICUM is hydrodispersable*. It is better dissolved in warm liquids. It can be mixed with either water, milk, or milk replacer, being slightly more soluble in water.



How to administer it through a stomach tube

Mix 50 g of RECOFAST ENTERICUM with 100 ml of warm liquid (water, milk or milk replacer).

*Dissolve a 100g sachet in 100ml-1L of liquid.

